


GCSE Mathematics (Grade 9-1)

Problem Solving – Sample 1 (Solutions)

H




fluidmaths.co.uk

{Aimed at students working towards a Grade 9 or 8}

The Grades and Marks shown are for guidance purposes only

The questions are repeated here for your convenience

- 1 An equilateral triangle of side 6cm is drawn inside a circle of area 108π
The triangle is drawn such that its height lies on the diameter of the circle as shown



Calculate the value of x exactly.

Solution

We can find x by calculating the difference between the length of the diameter of the circle and the length of the height of the triangle

$\sin 60 = \frac{h}{6}$ where h is the height of the triangle.

{Note that each angle in an equilateral triangle is 60° }

$$h = 6 \times \frac{\sqrt{3}}{2} = 3\sqrt{3} \quad \{\text{Note that exact value of } \sin 60 = \frac{\sqrt{3}}{2}\} \quad \text{[1mark]}$$

{Area of a circle = πr^2 }

Therefore, $108\pi = \pi r^2$

$$r^2 = 108$$

$$r = \sqrt{108} = 6\sqrt{3}$$

Hence, the length of the diameter will be $12\sqrt{3}$ [1mark]

The gaps between the base as well as the apex of the triangle and the circumference of the circle can be found as follows: $x + \frac{2}{3}x = \frac{5}{3}x$ [1mark]

Difference between the diameter of the circle and height of the triangle

$$12\sqrt{3} - 3\sqrt{3} = 9\sqrt{3}$$

$$\text{Therefore, } \frac{5}{3}x = 9\sqrt{3} \quad \text{[1mark]}$$

$$5x = 27\sqrt{3}$$

$$x = \frac{27\sqrt{3}}{5} \quad \text{[1mark]}$$

- 2 Given that, $5 : x(x + 4) = 3 : 7$
Show that $3x^2 + 12x - 35 = 0$

Solution

Write the ratio into a proportional equation

$$\frac{5}{x(x+4)} = \frac{3}{7} \quad \{\text{Cross multiply}\} \quad \text{[1mark]}$$

$$3x(x + 4) = 35 \quad \{\text{Expand the brackets}\}$$

$$3x^2 + 12x = 35 \quad \{\text{Subtract 35 from both sides}\} \quad \text{[1mark]}$$

$$\text{Hence, } 3x^2 + 12x - 35 = 0 \quad \text{[1mark]}$$

- 3 Given that $(\sqrt{p} + \sqrt{5p})^2 = 10 + q\sqrt{5}$
Show that $q = 3\frac{1}{3}$

Solution

Expand the brackets on the LHS

$$(\sqrt{p} + \sqrt{5p})^2 = (\sqrt{p} + \sqrt{5p})(\sqrt{p} + \sqrt{5p}) = p + p\sqrt{5} + p\sqrt{5} + 5p$$

{Simplify by collecting the like terms} [1mark]

$$p + p\sqrt{5} + p\sqrt{5} + 5p = 6p + 2p\sqrt{5} \quad \text{[1mark]}$$

{Now compare this expression with the expression on the RHS}

We notice that $6p = 10$

$$\text{Therefore, } p = \frac{10}{6} = \frac{5}{3} \quad \text{[1mark]}$$

Also, $q = 2p$

$$\text{Therefore, } q = 2 \times \frac{5}{3} = \frac{10}{3} = 3\frac{1}{3} \quad \text{[1mark]}$$

4

The solution for a quadratic equation is given as $x = \frac{-7 \pm \sqrt{49-24}}{6}$
 The coordinate $(3, k)$ lies on the curve. Find the exact value of k

Solution

We need to find the quadratic equation with such a solution

If $ax^2 + bx + c = 0$ then, $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

Now compare the quadratic formula above with the given value of x in the question: $x = \frac{-7 \pm \sqrt{49-24}}{6}$

Notice that,

$$-b = -7 \text{ Therefore, } b = 7$$

$$2a = 6 \text{ Therefore, } a = 3$$

$4ac = 24$ Substitute $a = 3$ into this equation to find the value of c

$$4 \times 3 \times c = 24 \text{ Therefore, } c = 2$$

[2marks]

Hence the quadratic equation is $3x^2 + 7x + 2 = 0$ **[1mark]**

Now, to find the value of k , we need to substitute $x = 3$ into the above equation

$$\text{If } x = 3 \text{ then, } 3(3)^2 + 7(3) + 2 = 0 \quad \textbf{[1mark]}$$

$$27 + 21 + 2 = 50$$

Therefore, $k = 50$ **[1mark]**

5 Given that $10x^2 - 9xy + 2y^2 = 0$

Find the possible values of $\frac{x}{y}$

Solution

We need to factorise the equation into a double bracket

We will then have to solve the factors in terms of x and y

The quadratic we have in this equation looks quite weird

However, it still factorises in a similar way as other quadratics

$$10x^2 - 9xy + 2y^2 = 0$$

Factorise into a double bracket as follows:

$$10 \times 2 = 20 \quad \{\text{Multiplying the coefficients of } x^2 \text{ and } y^2\}$$

We now need to find the factors of +20 which we can add to get -9

Therefore, both factors must be negative

The required factors will be -5 and -4

$$\text{Therefore, } 10x^2 - 9xy + 2y^2 = (10x - 5y)(10x - 4y)$$

{Now simplify both brackets}

$$(2x - y)(5x - 2y) = 0$$

[2marks]

Now solve each factor separately

$$\text{Either } 2x - y = 0$$

Therefore, $2x = y$ {Divide both sides by y and by 2}

$$\text{So we have } \frac{x}{y} = \frac{1}{2}$$

[1mark]

Or

$$5x - 2y = 0$$

$5x = 2y$ {Divide both sides by y and by 5}

$$\frac{x}{y} = \frac{2}{5}$$

[1mark]